Missouri Botanical Garden

Peter H. Raven Library

George Engelmann Botanical Notebooks



Pagination Note:
Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

Quercus sp.
foliis subintegris, cinereis
Aiken, S Carol, Ravenel

June 17 1877

×4

top of acorn

b

b

hovules
scattered, somewhat
removed from top

d

acorn depressed

Smithsonian Institution
U.S. National Herbarium
Washington, D.C. 20560 U.S.A.
Botanical Museum Leaflets
Vol. 28, No. 1, pp. 1-70
1980

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Deutsche Apotheke.

GEO. UDE,

Apothecary and Druggist

No. 3610 North 10th Street,

Between Mallinckrodt and Salisbury, **ST. LOUIS, MO.**

No. *187*

For .....

Rp.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

6034

0 cm 1 2 3 4 5 6 7 8 9 10
copyright reserved
MISSOURI BOTANICAL GARDEN

Dec 27 1876

G. apiculata
Mrs Smith. Oakland, June 15th

x10





Herbarium
Botanical Museum
Harvard University
Cambridge, Massachusetts

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

6035

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Leitch & Corlies
DRUGGISTS
South West Corner Fifth and Market Streets.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

G. corrinea Fresh, Lafayette Ar. July 3. 1876





white silk filling
The cells

first of last year blooming



Fertile ovule or young seed x10



sterile ovule



sterile ovules of different sizes all larger than the ovule of flower

two abortive ovules seen from back x10 or from outer side of cell

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Leitch & Corlies

DRUGGISTS

South West Corner Fifth and Market Streets.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

6036

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

G. coccinea (fresh)

July 3d 1876

very fresh 2 months old
×2

Cup tomentose
petiole
smooth green

green & smooth
no ovules seen



0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

6037

MISSOURI BOTANICAL GARDEN
ENGELMANN PAPERS

Leitch & Corlies
COMMISSION
DRUGGISTS
South West Corner Fifth and Market Streets.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Q. macrocarpa Lafayette Ave. July 4 1876

young fruit 2 months old

×2 ×2 ×2



sect



sect

some wool-like cells



×10 middle section

lower section

3 ribbons 3-celled above 1-celled

upper section



stigmata correspond with cells, therefore 3 pellagy

seeds (ovules) 2 months old almost equal yet in size and all fresh & green

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

6038

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Leitch & Corlies

IMPORTERS & JOBBERS

South West Corner Fifth and Market Streets.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Quercus alba

Aug 12 1876

at Judge Woerner's house near Arsenal

Cup full grown, acorns short (yet full length?) but apparently full formed & enlarged. Pericarp & kernel and its membranes quite soft yet and cells and dissepiments distinct.

×10

×30



Plants long hairy

Lowest part of fertile cell silky and the two cells with abortive ovules

hairy cell walls quite distant

silky

ovule erect!

×10

a

horizontal section through base of acorn; a is the coating of the seed (the membrane only) below cotyledons — 2 cells with 2 seed are open — others not touched, cells hairy

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

6039

Leitch & Conlies

DRUGGISTS

South West Corner Fifth and Market Streets.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Quamoclit coccinea — St Louis — Aug 13 1876

full size
Aug 13!
knobs of cupula scarcely visible yet at lowest part
cotyledons already firm except lowest part

×30

The whole county of the mid — yet sticky



×10
5 abortive ovules

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

6040

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Leitch & Corlies

WHOLESALE DRUGGISTS

South West Corner Fifth and Market Streets.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Jun 1868

Corylus & Carpinus in Petersburg



Castanea 4 spring
values of fruit, including
3 to 7 of aborted
flowers.



Fagus 4 spring
values of fruit, including
2 (central one aborted)
flowers



Quercus
circular, hemispherical
Cupula including a
single flower

over

The cupula of Quercus and the
spiny valves of Fagus & Castanea
correspond, are organs within
the last bracts, include an flower
or an inflorescence, and are
as as yet unexplained organs.
Scales of cupula of Quercus
without regular arrangement — no
leaves.

Some Quercus (Cyclobalanus)
have several verticils of Cupulae



MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

1408

over

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Dec 1. 1876

Cavities on cups of oaks, secondary abortive
fem flowers? Or made by insects?

Cavity seen in Q. bicolor, Kimmswick Sept 1858
Q. Prinus shore of Potomac Va; near Ebbitt, Ward &
Q. Castanea shady bottom woods opp Ft Sept 1860
Q. prinoides Olney Sept 28 1871

Cavity & acornlet in Q. prinoides Olney RI. Sept 28 1871

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

6042

spines: flowers short; seeds as in the last. — Pilocereus.

C. Schottii, Engelm., 4 to 10 feet high, the lower part 5-angled, ~~with distant~~ areolæ and few very short and stout spines; the upper flowering portion deeply 5-ribbed, with close-set areolæ bearing numerous setaceous spines, almost hiding the small flowers and small berries, — from the same localities as the two last species, may also be found in southern California.

4. Opuntia, Tourn., Miller.

Tube of the flower very short, cupshaped; petals spreading or rarely erect. Ovary ~~covered~~ with bristle-bearing areolæ in the axils of small terete deciduous sepals. ~~Fruit~~ Berry succulent or sometimes dry, marked with bristly or spiny areolæ, truncate with a wide umbilicus. Seeds large, white, compressed, ~~seeds~~ with the embryo coiled around the albumen: cotyledons large, foliaceous, —

Articulated much branched plants, of various shapes, low and prostrate, or erect and shrub like; young branches ~~bearing~~ with small terete subulate early deciduous leaves, and in their

MISSOURI BOTANICAL GARDEN ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

April 1876

Stamens in Oaks

White oaks with annual fruit — stamens 5–8

rarely (forms of undulata and of virens sometimes) only 4

very rarely (Michauxii) 10

White oaks with biennial fruit

Chrysolepis (5–) 10

Aegilops (Camp. Asia) 4–6

rarely (ovata) 2–4 2–4

Black oaks with annual fruit

Agrifolia 6–8 (–10)

pumila; hypoleuca 4

Black oaks with biennial fruit

2–5 (–6)

rarely more 4

Chestnut oaks (discolor) 10

Lepidobalanus filaments not more than twice as long as anthers (anthers mostly oval)

Cyclobalanopsis filaments many times longer than small globose anthers

---

All Official Letters to this Office should be addressed to THE

6043

and II. Botany. Captain Simpson's report of Explorations across the Great Basin of Utah. Please return them at as early a day as practicable.

By direction of the Chief of Engineers,

Very respectfully,

Your Obedient Servant,

Geo. H. Elliot.

Major of Engineers.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Jan 16th 1876

Nuces Quercus

Subsubsection

1. Nuces tenues intus glabrae lucentes

ann Q. alba, macrocarpa, Prinos, stellata
virens undulata,

biem Q. Suber, Libani, Pseudosuber (one gland
of this found loose in herb, given by J. Gay, was
silky tomentose, the others smooth.)

(Q. lyrata is similar but has a very thick
shell.)

3. Nuces crassiores intus cerrico tomentosae

biem Q. Toxodonii, rubra, nigra, palustris
imbricaria

ann Q. pumila, agrifolia

2. Nuces tenuiusculae (intus pubescentes)

biem Q. chrysolepis

4. Nuces crassissimae intus cerrico tomentosae

biem Q. densiflora

Knobs are peculiar to the cupulae
of the white oaks — the others have
flat scales

6044

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

St. Louis Jan 15th

Dr Geo. J. Engelmann!

Dear Sir!

You have omitted to put down the quantities of two articles in the Prescription for Mrs Scheel, which I sent you by bearer.

Please correct

and I remain

very respectfully

yours

H. Broste

for

Poole & Co.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN